arXiv:math/0509003v3 [math.GT] 23 Sep 2008

# HOMOTOPY ON SPATIAL GRAPHS AND THE SATO-LEVINE INVARIANT

THOMAS FLEMING AND RYO NIKKUNI

Abstract. Edge-homotopy and vertex-homotopy are equivalence relations on spatial graphs which are generalizations of Milnor's link-homotopy. We introduce some edge (resp. vertex)-homotopy invariants of spatial graphs by applying the Sato-Levine invariant for the 2-component constituent algebraically split links and show examples of non-splittable spatial graphs up to edge (resp. vertex)-homotopy, all of whose constituent links are link-homotopically trivial.

## 1. Introduction

Throughout this paper we work in the piecewise linear category. Let $G$ be a finite graph which does not have isolated vertices and free vertices. An embedding $f$ of $G$ into the 3-sphere $S^3$ is called a *spatial embedding of* $G$ or simply a *spatial graph*. For a spatial embedding $f$ and a subgraph $H$ of $G$ which is homeomorphic to the 1-sphere $S^1$ or a disjoint union of 1-spheres, we call $f(H)$ a *constituent knot* or a *constituent link* of $f$, respectively. A graph $G$ is said to be *planar* if there exists an embedding of $G$ into the 2-sphere $S^2$, and a spatial embedding of a planar graph is said to be *trivial* if it is ambient isotopic to an embedding of the graph into a 2-sphere in $S^3$. A spatial embedding $f$ of a graph $G$ is said to be *split* if there exists a 2-sphere $S$ in $S^3$ such that $S \cap f(G) = \emptyset$ and each component of $S^3 - S$ has intersection with $f(G)$, and otherwise $f$ is said to be *non-splittable*.

Two spatial embeddings of a graph $G$ are said to be *edge-homotopic* if they are transformed into each other by *self crossing changes* and ambient isotopies, where a self crossing change is a crossing change on the same spatial edge, and *vertex-homotopic* if they are transformed into each other by crossing changes on two adjacent spatial edges and ambient isotopies.[1] These equivalence relations were introduced by Taniyama [19] as generalizations of Milnor's *link-homotopy* on links [8], namely if $G$ is homeomorphic to a disjoint union of 1-spheres, then these are none other than link-homotopy. There are many studies about link-homotopy. In particular, the link-homotopy classification was given for 2- and 3-component links by Milnor [8], for 4-component links by Levine [7] and for all links by Habegger and

---

2000 *Mathematics Subject Classification.* Primary 57M15; Secondary 57M25.

*Key words and phrases.* Spatial graph, edge-homotopy, vertex-homotopy, Sato-Levine invariant.

The first author was supported by a Fellowship of the Japan Society for the Promotion of Science for Post-Doctoral Foreign Researchers (Short-Term) (No. PE05003).

The second author was partially supported by a Grant-in-Aid for Scientific Research (B) (2) (No. 15340019), Japan Society for the Promotion of Science.

[1] In [19], edge-homotopy and vertex-homotopy were called *homotopy* and *weak homotopy*, respectively.

Lin [2]. On the other hand, there are very few studies about edge (resp. vertex)-homotopy on spatial graphs [18], [9], [13], [11].

In [18], Taniyama defined an edge (resp. vertex)-homotopy invariant of spatial graphs called the *$\alpha$-invariant* by applying the *Casson invariant* (or equivalently the second coefficient of the *Conway polynomial*) of the constituent knots and showed that there exists a non-trivial spatial embedding $f$ of a planar graph up to edge (resp. vertex)-homotopy, even in the case where $f$ does not contain any constituent link. But the $\alpha$-invariant cannot detect a non-splittable spatial embedding of a disconnected graph up to edge (resp. vertex)-homotopy. As far as the authors know, an example of a non-splittable spatial embedding of a disconnected graph up to edge (resp. vertex)-homotopy, all of whose constituent links are link-homotopically trivial has not yet been demonstrated.

Our purpose in this paper is to study spatial embeddings of disconnected graphs up to edge (resp. vertex)-homotopy by applying the *Sato-Levine invariant* [14] (or equivalently the third coefficient of the Conway polynomial) for the constituent 2-component algebraically split links and show that there exist infinitely many non-splittable spatial embeddings of a certain disconnected graph up to edge (resp. vertex)-homotopy all of whose constituent links are link-homotopically trivial. These examples show that edge (resp. vertex)-homotopy on spatial graphs behaves quite differently than link-homotopy on links. In the next section we give the definitions of our invariants and state their invariance up to edge (resp. vertex)-homotopy.

## 2. Definitions of invariants

We call a subgraph of a graph $G$ a *cycle* if it is homeomorphic to the 1-sphere, and a cycle is called a *$k$-cycle* if it contains exactly $k$ edges. For a subgraph $H$ of $G$, we denote the set of all cycles of $H$ by $\Gamma(H)$. We set $\mathbb{Z}_m = \{0, 1, \ldots, m-1\}$ for a positive integer $m$ and $\mathbb{Z}_0 = \mathbb{Z}$. We regard $\mathbb{Z}_m$ as an abelian group in the obvious way. We call a map $\omega : \Gamma(H) \to \mathbb{Z}_m$ a *weight on* $\Gamma(H)$ *over* $\mathbb{Z}_m$. For an edge $e$ of $H$, we denote the set of all cycles of $H$ which contain the edge $e$ by $\Gamma_e(H)$. For a pair of two adjacent edges $e_1$ and $e_2$ of $H$, we denote the set of all cycles of $H$ which contain the edges $e_1$ and $e_2$ by $\Gamma_{e_1,e_2}(H)$. Then we say that a weight $\omega$ on $\Gamma(H)$ over $\mathbb{Z}_m$ is *weakly balanced* [2] *on an edge* $e$ if

$$\sum_{\gamma \in \Gamma_e(H)} \omega(\gamma) = 0$$

in $\mathbb{Z}_m$ [10], and *weakly balanced on a pair of adjacent edges* $e_1$ *and* $e_2$ if

$$\sum_{\gamma \in \Gamma_{e_1,e_2}(H)} \omega(\gamma) = 0$$

in $\mathbb{Z}_m$. Let $G = G_1 \cup G_2$ be a disjoint union of two connected graphs and $\omega_i : \Gamma(G_i) \to \mathbb{Z}_m$ a weight on $\Gamma(G_i)$ over $\mathbb{Z}_m$ $(i = 1, 2)$. Let $f$ be a spatial embedding of $G$ such that

$$\omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(f(\gamma), f(\gamma')) = 0$$

in $\mathbb{Z}$ for any $\gamma \in \Gamma(G_1)$ and $\gamma' \in \Gamma(G_2)$, where $\mathrm{lk}(L) = \mathrm{lk}(K_1, K_2)$ denotes the *linking number* of a 2-component oriented link $L = K_1 \cup K_2$. Then we define

---

[2] A weight $\omega$ on $\Gamma(H)$ over $\mathbb{Z}_m$ is said to be *balanced on an edge $e$ of $H$* if $\sum_{\gamma \in \Gamma_e(H)} \omega(\gamma)[\gamma] = 0$ in $H_1(H; \mathbb{Z}_m)$, where the orientation of $\gamma$ is induced by the one of $e$ [18].

$\beta_{\omega_1,\omega_2}(f) \in \mathbb{Z}_m$ by

$$\beta_{\omega_1,\omega_2}(f) \equiv \sum_{\substack{\gamma\in\Gamma(G_1)\\ \gamma'\in\Gamma(G_2)}} \omega_1(\gamma)\omega_2(\gamma')a_3(f(\gamma), f(\gamma')) \pmod{m},$$

where $a_3(L) = a_3(K_1, K_2)$ denotes the third coefficient of the Conway polynomial of a 2-component oriented link $L = K_1 \cup K_2$. We remark here that $a_3(L)$ coincides with the *Sato-Levine invariant* $\beta(L)$ of $L$ if $L$ is *algebraically split*, namely $\mathrm{lk}(K_1, K_2) = 0$ [1], [17]. Thus our $\beta_{\omega_1,\omega_2}(f)$ is also the modulo $m$ reduction of the summation of Sato-Levine invariants for the constituent 2-component algebraically split links of $f$.

**Remark 2.1.** For a 2-component algebraically split link $L = K_1 \cup K_2$,

(1) The value of $a_3(L)$ does not depend on the orientations of $K_1$ and $K_2$. Actually we can check it easily by the original definition of the Sato-Levine invariant.

(2) The value of $a_3(L)$ is not a link-homotopy invariant of $L$ (see also Lemma 3.1). For example, the Whitehead link $L$ is link-homotopically trivial but $a_3(L) = 1$.

Now we state the invariance of $\beta_{\omega_1,\omega_2}$ up to edge (resp. vertex)-homotopy under some conditions on the graphs.

**Theorem 2.2.** *Let $G = G_1 \cup G_2$ be a disjoint union of two connected graphs and $\omega_i$ a weight on $\Gamma(G_i)$ over $\mathbb{Z}_m$ $(i = 1, 2)$. Let $f$ be a spatial embedding of $G$ such that*

$$\omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(f(\gamma), f(\gamma')) = 0$$

*in $\mathbb{Z}$ for any $\gamma \in \Gamma(G_1)$ and $\gamma' \in \Gamma(G_2)$. Then we have the following:*

(1) *If $\omega_i$ is weakly balanced on any edge of $G_i$ $(i = 1, 2)$, then $\beta_{\omega_1,\omega_2}(f)$ is an edge-homotopy invariant of $f$.*

(2) *If $\omega_i$ is weakly balanced on any pair of adjacent edges of $G_i$ $(i = 1, 2)$, then $\beta_{\omega_1,\omega_2}(f)$ is a vertex-homotopy invariant of $f$.*

We prove Theorem 2.2 in the next section. In addition, by using an integer-valued invariant (Theorem 4.2), we show that there exist infinitely many non-splittable spatial embeddings of a certain disconnected graph up to edge-homotopy all of whose constituent links are link-homotopically trivial (Example 4.3). We also exhibit an infinite family of non-splittable spatial embeddings of a certain disconnected graph up to vertex-homotopy which can be distinguished by our integer-valued invariant (Example 4.4).

We note that if a graph $G$ contains a connected component which is homeomorphic to the 1-sphere, then our invariants in Theorem 2.2 are useless. For such cases, we can define edge (vertex)-homotopy invariants that take values in $\mathbb{Z}_2$ on weaker condition for weights than the one stated in Theorem 2.2. For a subgraph $H$ of a graph $G$, we say that a weight $\omega$ on $\Gamma(H)$ over $\mathbb{Z}_2$ is *totally balanced* if

$$\sum_{\gamma\in\Gamma(H)} \omega(\gamma)[\gamma] = 0$$

in $H_1(H; \mathbb{Z}_2)$. We note that if a weight $\omega$ on $\Gamma(H)$ over $\mathbb{Z}_2$ is totally balanced, then it is weakly balanced on any edge $e$ of $H$ (Lemma 3.2), but not always weakly balanced on any pair of adjacent edges of $H$ (Remark 3.3). Then we have the following.

**Theorem 2.3.** *Let* $G = G_1 \cup G_2$ *be a disjoint union of two connected graphs and* $\omega_i$ *a weight on* $\Gamma(G_i)$ *over* $\mathbb{Z}_2$ $(i = 1, 2)$. *Let* $f$ *be a spatial embedding of* $G$ *such that*

$$\omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(f(\gamma), f(\gamma')) = 0$$

*in* $\mathbb{Z}$ *for any* $\gamma \in \Gamma(G_1)$ *and* $\gamma' \in \Gamma(G_2)$. *Then we have the following:*

(1) *If either* $\omega_1$ *is totally balanced on* $\Gamma(G_1)$ *or* $\omega_2$ *is totally balanced on* $\Gamma(G_2)$, *then* $\beta_{\omega_1,\omega_2}(f)$ *is an edge-homotopy invariant of* $f$.

(2) *If either* $\omega_1$ *is totally balanced on* $\Gamma(G_1)$ *and weakly balanced on any pair of adjacent edges of* $G_1$, *or* $\omega_2$ *is totally balanced on* $\Gamma(G_2)$ *and weakly balanced on any pair of adjacent edges of* $G_2$, *then* $\beta_{\omega_1,\omega_2}(f)$ *is a vertex-homotopy invariant of* $f$.

We also prove Theorem 2.3 in the next section and give some examples in Section 5. In particular, we show that there exist infinitely many non-splittable spatial embeddings of a certain disconnected graph up to vertex-homotopy, all of whose constituent links are link-homotopically trivial (Example 5.4). We remark here that the $\mathbb{Z}_2$-valued invariant in Theorem 2.3 cannot always be extended to an integer-valued one (Remark 5.5).

Theorems 2.2 and 2.3 do not work for spatial graphs as illustrated in Figure 2.1, for instance. In Section 6, we state a method to detect such non-splittable spatial graphs up to edge-homotopy by using a planar surface having a graph as a spine (Theorem 6.1). Actually we show that each of the spatial graphs as illustrated in Figure 2.1 is non-splittable up to edge-homotopy (Example 6.2).



FIGURE 2.1.

## 3. PROOFS OF THEOREMS 2.2 AND 2.3

We first calculate the change in the third coefficient of the Conway polynomial of 2-component algebraically split links which differ by a single self crossing change.

**Lemma 3.1.** *Let* $L_+$ *and* $L_-$ *be two* 2*-component oriented links and* $L_0 = J_1 \cup J_2 \cup K$ *a* 3*-component oriented link which are identical except inside the depicted regions as illustrated in Figure 3.1. Suppose that* $\mathrm{lk}(L_+) = \mathrm{lk}(L_-) = 0$. *Then it holds that*

$$a_3(L_+) - a_3(L_-) = -\mathrm{lk}(J_1, K)^2 = -\mathrm{lk}(J_2, K)^2.$$



FIGURE 3.1.

*Proof.* By the skein relation of the Conway polynomial and a well-known formula for the second coefficient of the Conway polynomial of a 3-component oriented link (cf. [4], [3], [5]), we have that

$$\begin{aligned} a_3(L_+) - a_3(L_-) &= \mathrm{lk}(J_1, J_2)\mathrm{lk}(J_2, K) + \mathrm{lk}(J_2, K)\mathrm{lk}(J_1, K) \\ &\quad + \mathrm{lk}(J_1, K)\mathrm{lk}(J_1, J_2). \end{aligned} \tag{3.1}$$

We note that

$$\mathrm{lk}(J_1, K) + \mathrm{lk}(J_2, K) = 0 \tag{3.2}$$

by the condition $\mathrm{lk}(L_+) = \mathrm{lk}(L_-) = 0$. Thus by (3.1) and (3.2), we have that

$$\begin{aligned} a_3(L_+) - a_3(L_-) &= \mathrm{lk}(J_1, J_2)\{-\mathrm{lk}(J_1, K)\} + \mathrm{lk}(J_2, K)\mathrm{lk}(J_1, K) \\ &\quad + \mathrm{lk}(J_1, K)\mathrm{lk}(J_1, J_2) \\ &= \mathrm{lk}(J_2, K)\mathrm{lk}(J_1, K). \end{aligned}$$

Therefore by (3.2) we have the result. □

*Proof of Theorem 2.2.* (1) Let $f$ and $g$ be two spatial embeddings of $G$ such that

$$\omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(f(\gamma), f(\gamma')) = 0 \tag{3.3}$$

in $\mathbb{Z}$ for any $\gamma \in \Gamma(G_1)$ and $\gamma' \in \Gamma(G_2)$ and $g$ is edge-homotopic to $f$. Then it also holds that

$$\omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(g(\gamma), g(\gamma')) = 0 \tag{3.4}$$

in $\mathbb{Z}$ for any $\gamma \in \Gamma(G_1)$ and $\gamma' \in \Gamma(G_2)$ because the linking number of a 2-component constituent link of a spatial graph is an edge-homotopy invariant. First we show that if $f$ is transformed into $g$ by self crossing changes on $f(G_1)$ and ambient isotopies, then $\beta_{\omega_1,\omega_2}(f) = \beta_{\omega_1,\omega_2}(g)$. It is clear that any link invariant of a constituent link of a spatial graph is also an ambient isotopy invariant of the spatial graph. Thus we may assume that $g$ is obtained from $f$ by a single crossing change on $f(e)$ for an edge $e$ of $G_1$ as illustrated in Figure 3.2. Moreover, by smoothing this crossing point we can obtain the spatial embedding $h$ of $G$ and the knot $J_h$ as illustrated in Figure 3.2. Then by (3.3), (3.4), Lemma 3.1 and the assumption for $\omega_1$ we have that

$$\begin{aligned}
\beta_{\omega_1,\omega_2}(f)-\beta_{\omega_1,\omega_2}(g) &\equiv \sum_{\substack{\gamma\in\Gamma(G_1)\\ \gamma'\in\Gamma(G_2)}} \omega_1(\gamma)\omega_2(\gamma')\left\{a_3(f(\gamma),f(\gamma'))-a_3(g(\gamma),g(\gamma'))\right\}\\
&= \sum_{\substack{\gamma\in\Gamma_e(G_1)\\ \gamma'\in\Gamma(G_2)}} \omega_1(\gamma)\omega_2(\gamma')\left\{a_3(f(\gamma),f(\gamma'))-a_3(g(\gamma),g(\gamma'))\right\}\\
&= -\sum_{\substack{\gamma\in\Gamma_e(G_1)\\ \gamma'\in\Gamma(G_2)}} \omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(h(\gamma'),J_h)^2\\
&= -\left(\sum_{\gamma\in\Gamma_e(G_1)}\omega_1(\gamma)\right)\sum_{\gamma'\in\Gamma(G_2)}\omega_2(\gamma')\mathrm{lk}(h(\gamma'),J_h)^2\\
&\equiv 0.
\end{aligned}$$

Therefore we have that $\beta_{\omega_1,\omega_2}(f)=\beta_{\omega_1,\omega_2}(g)$. In the same way we can show that if $f$ is transformed into $g$ by self crossing changes on $f(G_2)$ and ambient isotopies, then $\beta_{\omega_1,\omega_2}(f)=\beta_{\omega_1,\omega_2}(g)$. Thus we have that $\beta_{\omega_1,\omega_2}$ is an edge-homotopy invariant.

(2) By considering the triple of spatial embeddings as illustrated in Figure 3.3, we can prove (2) in a similar way as the proof of (1). We omit the details. ☐



FIGURE 3.2.

Next we prove Theorem 2.3. For a subgraph $H$ of a graph $G$, we have the following.

**Lemma 3.2.** *A totally balanced weight $\omega$ on $\Gamma(H)$ over $\mathbb{Z}_2$ is weakly balanced on any edge $e$ of $H$.*

*Proof.* For an edge $e$ of $H$, we can represent any $\gamma\in\Gamma_e(H)$ as $e+c_\gamma\in Z_1(H;\mathbb{Z}_2)$, where $c_\gamma$ is a 1-chain in $C_1(H\setminus e;\mathbb{Z}_2)$. Then we have that

$$\begin{aligned}
0 &= \sum_{\gamma\in\Gamma(H)}\omega(\gamma)[\gamma]\\
&= \sum_{\gamma\in\Gamma_e(H)}\omega(\gamma)[e+c_\gamma]+\sum_{\gamma'\in\Gamma(H)\setminus\Gamma_e(H)}\omega(\gamma')[\gamma']
\end{aligned}$$



Figure 3.3.

in $H_1(H;\mathbb{Z}_2)$. This implies that if $\omega$ is not weakly balanced on $e$, then $\omega$ is not totally balanced on $\Gamma(H)$ over $\mathbb{Z}_2$. □

**Remark 3.3.** A totally balanced weight $\omega$ on $\Gamma(H)$ over $\mathbb{Z}_2$ is not always weakly balanced on any pair of adjacent edges of $H$. For example, let $\omega$ be a weight on $\Theta_3$ (see Example 4.3) over $\mathbb{Z}_2$ defined by $\omega(\gamma)=1$ for any cycle $\gamma\in\Gamma(\Theta_3)$. It is easy to see that $\omega$ is totally balanced, but not weakly balanced, on each pair of adjacent edges of $\Theta_3$.

*Proof of Theorem 2.3.* (1) Let $f$ and $g$ be two spatial embeddings of $G$ which are edge-homotopic such that

$$\omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(f(\gamma),f(\gamma'))=\omega_1(\gamma)\omega_2(\gamma')\mathrm{lk}(g(\gamma),g(\gamma'))=0$$

in $\mathbb{Z}$ for any $\gamma\in\Gamma(G_1)$ and $\gamma'\in\Gamma(G_2)$. First we show that if $f$ is transformed into $g$ by self crossing changes on $f(G_1)$ and ambient isotopies, then $\beta_{\omega_1,\omega_2}(f)=\beta_{\omega_1,\omega_2}(g)$. In the same way as the proof of Theorem 2.2, we may consider three spatial embeddings $f,g$ and $h$ of $G$ and the knot $J_h$ as illustrated in Figure 3.2. Then, by the same calculation in the proof of Theorem 2.2, we have that

$$\begin{aligned}\beta_{\omega_1,\omega_2}(f)-\beta_{\omega_1,\omega_2}(g) &\equiv -\left(\sum_{\gamma\in\Gamma_e(G_1)}\omega_1(\gamma)\right)\sum_{\gamma'\in\Gamma(G_2)}\omega_2(\gamma')\mathrm{lk}(h(\gamma'),J_h)^2\\ &\equiv \left(\sum_{\gamma\in\Gamma_e(G_1)}\omega_1(\gamma)\right)\sum_{\gamma'\in\Gamma(G_2)}\omega_2(\gamma')\mathrm{lk}(h(\gamma'),J_h)\\ &\equiv \left(\sum_{\gamma\in\Gamma_e(G_1)}\omega_1(\gamma)\right)\mathrm{lk}\left(\sum_{\gamma'\in\Gamma(G_2)}\omega_2(\gamma')h(\gamma'),J_h\right).\end{aligned}$$

If $\omega_1$ is totally balanced on $\Gamma(G_1)$, then by Lemma 3.2 it is weakly balanced on any edge $e$ of $G_1$. This implies that $\beta_{\omega_1,\omega_2}(f)=\beta_{\omega_1,\omega_2}(g)$. If $\omega_2$ is totally balanced on

$\Gamma(G_1)$, then we have that

$$\mathrm{lk}\left(\sum_{\gamma'\in\Gamma(G_2)}\omega_2(\gamma')h(\gamma'), J_h\right) \equiv \mathrm{lk}\,(0, J_h) = 0.$$

Therefore this also implies that $\beta_{\omega_1,\omega_2}(f) = \beta_{\omega_1,\omega_2}(g)$. In the same way we can show that if $f$ is transformed into $g$ by self crossing changes on $f(G_2)$ and ambient isotopies, then $\beta_{\omega_1,\omega_2}(f) = \beta_{\omega_1,\omega_2}(g)$. Thus we have that $\beta_{\omega_1,\omega_2}$ is an edge-homotopy invariant.

(2) By considering the triple of spatial embeddings as illustrated in Figure 3.3, we can prove (2) in a similar way as the proof of (1). We also omit the details. □

Since the Conway polynomial of a split link is zero, our invariants take the value zero for any split (2-component) spatial graph. Therefore if the value of our invariant of a spatial graph is not zero, then it is non-splittable up to edge (resp. vertex)-homotopy.

## 4. INTEGER-VALUED INVARIANTS

Let $G$ be a planar graph. An embedding $p : G \to S^2$ is said to be *cellular* if the closure of each of the connected components of $S^2 - p(G)$ is homeomorphic to the disk. Then we regard the set of the boundaries of all of the connected components of $S^2 - p(G)$ as a subset of $\Gamma(G)$ and denote it by $\Gamma_p(G)$. We say that $G$ *admits a checkerboard coloring on* $S^2$ if there exists a cellular embedding $p : G \to S^2$ such that we can color all of the connected components of $S^2 - p(G)$ by two colors (black and white) so that any of the two components which are adjacent by an edge have distinct colors; see Figure 4.1. We denote the subset of $\Gamma_p(G)$ which corresponds to the black (resp. white) colored components by $\Gamma^b_p(G)$ (resp. $\Gamma^w_p(G)$).



FIGURE 4.1.

**Proposition 4.1.** *Let $G$ be a planar graph which is not homeomorphic to $S^1$ and admits a checkerboard coloring on $S^2$ with respect to a cellular embedding $p : G \to S^2$. Let $\omega_p$ be a weight on $\Gamma(G)$ over $\mathbb{Z}$ defined by*

$$\omega_p(\gamma) = \begin{cases} 1 & (\gamma \in \Gamma^b_p(G)), \\ -1 & (\gamma \in \Gamma^w_p(G)), \\ 0 & (\gamma \in \Gamma(G) \setminus \Gamma_p(G)). \end{cases}$$

*Then $\omega_p$ is weakly balanced on any edge of $G$.*

*Proof.* For any edge $e$ of $G$, there exist exactly two cycles $\gamma \in \Gamma^b_p(G)$ and $\gamma' \in \Gamma^w_p(G)$ such that $e \subset \gamma$ and $e \subset \gamma'$. Thus we have the result. □

We call the weight $\omega_p$ in Proposition 4.1 a *checkerboard weight*. Thus by Proposition 4.1 and Theorem 2.2 (1), we can obtain an integer-valued edge-homotopy invariant as follows.

**Theorem 4.2.** *Let $G = G_1 \cup G_2$ be a disjoint union of two connected planar graphs such that $G_i$ is not homeomorphic to $S^1$ and admits a checkerboard coloring on $S^2$ with respect to a cellular embedding $p_i : G_i \to S^2$ $(i = 1, 2)$. Let $\omega_{p_i}$ be a checkerboard weight on $\Gamma(G_i)$ over $\mathbb{Z}$ $(i = 1, 2)$ and $f$ a spatial embedding of $G$ such that*

$$\omega_{p_1}(\gamma)\omega_{p_2}(\gamma')\mathrm{lk}(f(\gamma), f(\gamma')) = 0$$

*in $\mathbb{Z}$ for any $\gamma \in \Gamma(G_1)$ and $\gamma' \in \Gamma(G_2)$. Then $\beta_{\omega_{p_1},\omega_{p_2}}(f)$ is an integer-valued edge-homotopy invariant of $f$.* □

**Example 4.3.** Let $\Theta_n$ be a graph with two vertices $u$ and $v$ and $n$ edges $e_1, e_2, \ldots, e_n$, each of which joins $u$ and $v$. A spatial embedding of $\Theta_n$ is called a *(spatial) theta $n$-curve* or simply a *theta curve* if $n = 3$. For $n \geq 2$, we denote that a cycle of $\Theta_n$ consists of two edges $e_i$ and $e_j$ by $\gamma_{ij}$ $(i < j)$. Then it is clear that $\Theta_n$ admits a cellular embedding $p : \Theta_n \to S^2$ so that

$$\Gamma_p(\Theta_n) = \{\gamma_{12}, \gamma_{23}, \ldots, \gamma_{n-1,n}, \gamma_{1n}\}.$$

Moreover, for $m \geq 1$, $\Theta_{2m}$ admits a checkerboard coloring on $S^2$ so that

$$\begin{aligned}\Gamma_p^b(\Theta_{2m}) &= \{\gamma_{12}, \gamma_{34}, \ldots, \gamma_{2m-1,2m}\}, \\ \Gamma_p^w(\Theta_{2m}) &= \{\gamma_{23}, \gamma_{45}, \ldots, \gamma_{2m-2,2m-1}, \gamma_{1,2m}\}.\end{aligned}$$

Now let $G$ be a disjoint union of two copies of $\Theta_4$, each of which admits a checkerboard coloring on $S^2$ with respect to the cellular embedding $p$ as above. Let $\omega_p$ be a checkerboard weight on $\Gamma(\Theta_4)$ over $\mathbb{Z}$ and $g_1$ a spatial embedding of $G$ as illustrated in Figure 4.2. We can see that any of the 2-component constituent links of $g_1$ has a zero linking number. More precisely, $g_1$ contains exactly one non-trivial 2-component link $L = g_1(\gamma_{14}) \cup g_1(\gamma'_{14})$ whose linking number is zero. Thus by Theorem 4.2 we have that $\beta_{\omega_p,\omega_p}(g_1)$ is an integer-valued edge-homotopy invariant of $g_1$. Then, by a direct calculation we have that $a_3(L) = 2$, namely $\beta_{\omega_p,\omega_p}(g_1) = 2$. Note that a 2-component link is link-homotopically trivial if and only if its linking number is zero [8]. This implies that $g_1$ is non-splittable up to edge-homotopy despite the fact that any of the constituent links of $g_1$ is link-homotopically trivial.



FIGURE 4.2.

Moreover, for an integer $m$, let $g_m$ be a spatial embedding of $G$ as illustrated in Figure 4.3. If $m \neq 0$, we can see that $g_m$ contains exactly one non-trivial 2-component link $L = g_m(\gamma_{14}) \cup g_m(\gamma'_{14})$ whose linking number is zero. Thus we also have that $\beta_{\omega_p,\omega_p}(g_m)$ is an integer-valued edge-homotopy invariant of $g_m$. Then, by a calculation we have that $a_3(L) = 2m$, namely $\beta_{\omega_p,\omega_p}(g_m) = 2m$. This implies that there exist infinitely many non-splittable spatial embeddings of $G$ up to edge-homotopy, all of whose constituent links are link-homotopically trivial.



FIGURE 4.3.

**Example 4.4.** Let $H$ be a graph as illustrated in Figure 4.4. We denote the cycle of $H$ which contains $e_i$ and $e_j$ by $\gamma_{ij}$ $(i < j)$. Let $G$ be a disjoint union of two copies of $H$ and $g_1$ a spatial embedding of $G$ as illustrated in Figure 4.5. This spatial embedding $g_1$ contains exactly one 4-component constituent link $L = g_1(\gamma_{12} \cup \gamma_{34} \cup \gamma'_{12} \cup \gamma'_{34})$. Note that if $g_1$ is split up to vertex-homotopy, then $L$ is split up to link-homotopy. Since $|\mu_{1234}(L)| = 1$, where $\mu_{1234}$ denotes Milnor's *$\mu$-invariant* of length 4 of 4-component links [8], we have that $L$ is non-splittable up to link-homotopy. Therefore we have that $g_1$ is non-splittable up to vertex-homotopy.

We can also prove this fact by our integer-valued vertex-homotopy invariant as follows. Let $\omega$ be a weight on $\Gamma(H)$ over $\mathbb{Z}$ defined by $\omega(\gamma_{14}) = \omega(\gamma_{23}) = 1$,

$\omega(\gamma_{13}) = \omega(\gamma_{24}) = -1$ and $\omega(\gamma) = 0$ if $\gamma$ is a 2-cycle. Then it is easy to see that $\omega$ is weakly balanced on any pair of adjacent edges of $H$. We can see that $g_1$ contains exactly one non-trivial 2-component constituent link $M = g_1(\gamma_{14} \cup \gamma'_{14})$ with $\mathrm{lk}(M) = 0$ and $a_3(M) = 2$. Thus by Theorem 2.2 (2) we have that $\beta_{\omega,\omega}(g_1)$ is an integer-valued vertex-homotopy invariant of $g_1$ and $\beta_{\omega,\omega}(g_1) = 2$. This implies that $g_1$ is non-splittable up to vertex-homotopy.



Figure 4.4.



Figure 4.5.

Moreover, let $g_m$ be a spatial embedding of $G$ as illustrated in Figure 4.5, which can be constructed in the same way as in Example 4.3. Then we can see that $\beta_{\omega,\omega}(g_m)$ is an integer-valued vertex-homotopy invariant of $g_m$ and $\beta_{\omega,\omega}(g_m) = 2m$. This implies that $g_m$ is non-splittable up to vertex-homotopy for any integer $m \neq 0$ and $g_i$ and $g_j$ are not vertex-homotopic for any $i \neq j$.

## 5. Modulo two invariants

**Proposition 5.1.** *Let $G$ be a planar graph which is not homeomorphic to $S^1$ and $p : G \to S^2$ a cellular embedding. Let $\omega_p : \Gamma(G) \to \mathbb{Z}_2$ be a weight on $\Gamma(G)$ over $\mathbb{Z}_2$ defined by*

$$\omega_p(\gamma) = \begin{cases} 1 & (\gamma \in \Gamma_p(G)), \\ 0 & (\gamma \in \Gamma(G) \setminus \Gamma_p(G)). \end{cases}$$

*Then $\omega_p$ is totally balanced.*

*Proof.* It holds that

$$\sum_{\gamma \in \Gamma(G)} \omega_p(\gamma)[\gamma] = \sum_{\gamma \in \Gamma_p(G)} [\gamma] = 2\left[\sum_{e \in E(G)} e\right] = 0$$

in $H_1(G;\mathbb{Z}_2)$, where $E(G)$ denotes the set of all edges of $G$. Thus we have the result. $\square$

Thus by Proposition 5.1 and Theorem 2.3 (1), we can obtain an edge-homotopy invariant as follows.

**Theorem 5.2.** *Let* $G = G_1 \cup G_2$ *be a disjoint union of two connected graphs such that* $G_1$ *is planar, not homeomorphic to* $S^1$ *and admits a cellular embedding* $p_1 : G_1 \to S^2$. *Let* $\omega_{p_1}$ *be a weight on* $\Gamma(G_1)$ *over* $\mathbb{Z}_2$ *as in Proposition 5.1,* $\omega_2$ *a weight on* $\Gamma(G_2)$ *over* $\mathbb{Z}_2$ *and* $f$ *a spatial embedding of* $G$ *such that*

$$\omega_{p_1}(\gamma)\omega_2(\gamma')\mathrm{lk}(f(\gamma), f(\gamma')) = 0$$

*in* $\mathbb{Z}$ *for any* $\gamma \in \Gamma(G_1)$ *and* $\gamma' \in \Gamma(G_2)$. *Then* $\beta_{\omega_{p_1},\omega_2}(f)$ *is an edge-homotopy invariant of* $f$. $\square$

**Example 5.3.** Let $G$ be a disjoint union of $\Theta_3$ and a circle $\gamma$. Let $\omega_p$ be a weight on $\Gamma(\Theta_3)$ over $\mathbb{Z}_2$ as in Proposition 5.1 with respect to a cellular embedding $p : \Theta_3 \to S^2$ as in Example 4.3, and $\omega$ a weight on $\Gamma(\gamma)$ over $\mathbb{Z}_2$ defined by $\omega(\gamma) = 1$. Let $g$ be a spatial embedding of $G$ as illustrated in Figure 5.1 (1). We can see that $g$ contains exactly one non-trivial 2-component link $L = g(\gamma_{13}) \cup g(\gamma)$ which is the Whitehead link, so $\mathrm{lk}(L) = 0$ and $a_3(L) = 1$. Thus by Theorem 5.2 we have that $\beta_{\omega_p,\omega}(g)$ is an edge-homotopy invariant of $g$ and $\beta_{\omega_p,\omega}(g) = 1$. Namely $g$ is non-splittable up to edge-homotopy despite the fact that any of the constituent links of $g$ is link-homotopically trivial.

**Example 5.4.** Let $G$ be a disjoint union of the complete bipartite graph on $3+3$ vertices $K_{3,3}$ and a circle $\gamma$. Let $\omega_{3,3}$ be a weight on $K_{3,3}$ over $\mathbb{Z}_2$ defined by $\omega_{3,3}(\gamma') = 1$ if $\gamma'$ is a 4-cycle and 0 if $\gamma'$ is a 6-cycle. Let $\omega$ be a weight on $\Gamma(\gamma)$ over $\mathbb{Z}_2$ defined by $\omega(\gamma) = 1$. Then it is not hard to see that $\omega_{3,3}$ is totally balanced and weakly balanced on any pair of adjacent edges of $K_{3,3}$. For a positive integer $m$, let $g_m$ be a spatial embedding of $G$ as illustrated in Figure 5.1 (2). Note that $g_i(K_{3,3})$ and $g_j(K_{3,3})$ are not vertex-homotopic for any $i \neq j$ [9]; namely $g_i$ and $g_j$ are not vertex-homotopic for any $i \neq j$. Since all of the 2-component constituent links of $g_m$ are algebraically split, by Theorem 2.3 (2) we have that $\beta_{\omega_{3,3},\omega}(g)$ is a vertex-homotopy invariant of $g_m$. Moreover we can see that there exists exactly one 4-cycle $\gamma'$ of $K_{3,3}$ so that $L = g_m(\gamma \cup \gamma')$ is non-trivial. Since $L$ is the Whitehead link, we have that $\beta_{\omega_{3,3},\omega}(g_m) = 1$. Therefore $g_m$ is non-splittable up to vertex-homotopy despite the fact that any of the constituent links of $g$ is link-homotopically trivial.



(1) (2)

Figure 5.1.

**Remark 5.5.** The $\mathbb{Z}_2$-valued invariant in Theorem 2.3 cannot always be extended to an integer-valued one. For example,

(1) Let us consider the graph $G$ and the invariant $\beta_{\omega_p,\omega}$ as in Example 5.3. Let $f$ be a spatial embedding of $G$ as illustrated in Figure 5.2. We can see that $f$ is edge-homotopic to the trivial spatial embedding $h$ of $G$. But by a calculation we have that $\sum_{1\leq i<j\leq 3} a_3(f(\gamma_{ij}), f(\gamma)) = -2$.



FIGURE 5.2.

(2) Let $G$ be a disjoint union of $\Theta_4$ and a circle $\gamma$. Let $\omega_p$ be a checkerboard weight on $\Gamma(\Theta_4)$ over $\mathbb{Z}$ as in Example 4.3. Note that the modulo two reduction of a checkerboard weight is totally balanced. So by Theorem 2.3 (1), the modulo two reduction of $\sum_{\gamma_{ij}\in\Gamma(\Theta_4)} \omega_p(\gamma_{ij}) a_3(f(\gamma_{ij}\cup\gamma))$ is an edge-homotopy invariant of a spatial embedding $f$ of $G$. Moreover, we can see that the integer-value $\sum_{\gamma_{ij}\in\Gamma(\Theta_4)} \omega_p(\gamma_{ij}) a_3(f(\gamma_{ij}\cup\gamma))$ is invariant under the self crossing change on $f(\Theta_4)$ in the same way as in the proof of Theorem 2.2 (1). But this value may change under a self crossing change on $f(\gamma)$. For example, let $f$ and $g$ be two spatial embeddings of $G$ as illustrated in Figure 5.3. We can see that $f$ is edge-homotopic to $g$. But by a calculation we have that

$$\begin{aligned}\sum_{\gamma_{ij}\in\Gamma(\Theta_4)} \omega_p(\gamma_{ij}) a_3(f(\gamma_{ij}), f(\gamma)) &= -1,\\ \sum_{\gamma_{ij}\in\Gamma(\Theta_4)} \omega_p(\gamma_{ij}) a_3(g(\gamma_{ij}), g(\gamma)) &= 1.\end{aligned}$$



FIGURE 5.3.

## 6. Applying the boundary of a planar surface

Let $X$ be a disjoint union of a graph $G$ and a planar surface $F$ with boundary. Let $\omega$ be a weight on $\Gamma(G)$ over $\mathbb{Z}_2$ and $\varphi$ an embedding of $X$ into $S^3$ such that

$$\omega(\gamma)\mathrm{lk}(\varphi(\gamma),\varphi(\gamma'))=0$$

in $\mathbb{Z}$ for any $\gamma\in\Gamma(G)$ and $\gamma'\in\Gamma(\partial F)$. Then we define $\beta_\omega(\varphi)\in\mathbb{Z}_2$ by

$$\beta_\omega(\varphi)\equiv\sum_{\substack{\gamma\in\Gamma(G)\\ \gamma'\in\Gamma(\partial F)}}\omega(\gamma)a_3(\varphi(\gamma),\varphi(\gamma'))\pmod 2.$$

Let $G$ be a disjoint union of a connected graph $G_1$ and a connected planar graph $G_2$. Let $f$ be a spatial embedding of $G$ and $p$ an embedding of $G_2$ into $S^2$. We denote the regular neighborhood of $p(G_2)$ in $S^2$ by $F(G_2;p)$, which is a planar surface having $p(G_2)$ as a spine. Then the spatial embedding $f$ induces an embedding $\tilde{f}_p$ of the disjoint union $G_1\cup F(G_2;p)$ into $S^3$, so that $\tilde{f}_p(G_1)=f(G_1)$ and $\tilde{f}_p(F(G_2;p))$ has $f(G_2)$ as a spine in the natural way. Note that such an induced embedding $\tilde{f}_p$ is not unique up to ambient isotopy. Let $\omega$ be a weight on $\Gamma(G_1)$ over $\mathbb{Z}_2$ so that

$$\omega(\gamma)\mathrm{lk}(\tilde{f}_p(\gamma),\tilde{f}_p(\gamma'))=0$$

in $\mathbb{Z}$ for any $\gamma\in\Gamma(G_1)$ and $\gamma'\in\Gamma(\partial F(G_2;p))$. Then we have the following.

**Theorem 6.1.** *If $f$ is split up to edge-homotopy, then $\beta_\omega(\tilde{f}_p)=0$ for any induced embedding $\tilde{f}_p$ of $G_1\cup F(G_2;p)$.*

*Proof.* By the assumption we have that $f$ is transformed into a split spatial embedding $u$ of $G$ by self crossing changes and ambient isotopies. Then each of the self crossing changes induces a self crossing change on $\tilde{f}_p(G_1)$ or a *band-pass move* [6] (see Figure 6.1) on $\tilde{f}_p(F(G_2;p))$. Namely $\tilde{f}_p$ can be transformed into an induced embedding $\tilde{u}_p$ of $G_1\cup F(G_2;p)$ by such moves and ambient isotopies. Let $\tilde{g}_p$ be an embedding of $G_1\cup F(G_2;p)$ into $S^3$ obtained from $\tilde{f}_p$ by a single self crossing change on $\tilde{f}_p(G_1)$ or a single band-pass move on $\tilde{f}_p(F(G_2;p))$. Then it still holds that

$$\omega(\gamma)\mathrm{lk}(\tilde{g}_p(\gamma),\tilde{g}_p(\gamma'))=0$$

in $\mathbb{Z}$ for any $\gamma\in\Gamma(G_1)$ and $\gamma'\in\Gamma(\partial F(G_2;p))$.

**Claim.** $\beta_\omega(\tilde{f}_p)=\beta_\omega(\tilde{g}_p)$.

Assume that $\tilde{g}_p$ is obtained from $\tilde{f}_p$ by a single self crossing change on $\tilde{f}_p(G_1)$. Since it holds that

$$\sum_{\gamma'\in\Gamma(\partial F(G_2;p))}[\gamma']=0$$

in $H_1(F(G_2;p);\mathbb{Z}_2)$, we can see that $\beta_\omega(\tilde{f}_p)=\beta_\omega(\tilde{g}_p)$ in a similar way as the proof of Theorem 2.3 (1). Next we assume that $\tilde{g}_p$ is obtained from $\tilde{f}_p$ by a single band-pass move on $\tilde{f}_p(F(G_2;p))$. Then $\tilde{g}_p|_{G_1\cup\partial F(G_2;p)}$ is obtained from $\tilde{f}_p|_{G_1\cup\partial F(G_2;p)}$ by a single *pass move* [6] (see Figure 6.1) on $\tilde{f}_p(\partial F(G_2;p))$. We divide our situation into the following two cases.

**Case 1.** Four strings in the pass move belong to $\tilde{f}_p(\gamma_1')$ and $\tilde{f}_p(\gamma_2')$ for exactly two cycles $\gamma_1'$ and $\gamma_2'$ in $\Gamma(\partial F(G_2;p))$.

This pass move causes a single self crossing change on $\tilde{f}_p(\gamma_1')$ and a single self crossing change on $\tilde{f}_p(\gamma_2')$. Then the separated components that result from smoothing each of the self crossings are orientation-reversing parallel knots; see Figure 6.2. So the difference between $\beta_\omega(\tilde{f}_p)$ and $\beta_\omega(\tilde{g}_p)$ is cancelled out in a similar way as in the proof of Theorem 2.2 (1). Thus we have that $\beta_\omega(\tilde{f}_p) = \beta_\omega(\tilde{g}_p)$.

**Case 2.** Four strings in the pass move belong to $\tilde{f}_p(\gamma')$ for a cycle $\gamma'$ in $\Gamma(\partial F(G_2; p))$.

It is known that a pass move on the same component of a proper link $L = J_1 \cup J_2 \cup \cdots \cup J_n$ preserves $\overline{\mathrm{Arf}}(L) \equiv \mathrm{Arf}(L) - \sum_{i=1}^{n} \mathrm{Arf}(J_i) \in \mathbb{Z}_2$ (cf. [16]).[3] Especially, if $n = 2$ then $a_3(L) \equiv \overline{\mathrm{Arf}}(L) \pmod 2$ [12, Lemma 3.5 (ii)]. Therefore in this case the pass move preserves $\omega(\gamma)a_3(\tilde{f}_p(\gamma), \tilde{f}_p(\gamma'))$ for any cycle $\gamma \in \Gamma(G_1)$. This implies that $\beta_\omega(\tilde{f}_p) = \beta_\omega(\tilde{g}_p)$.

Now by the argument above, we have that $\beta_\omega(\tilde{f}_p) = \beta_\omega(\tilde{u}_p)$. Then, each 2-component link $\tilde{u}_p(\gamma \cup \gamma')$ is split for any $\gamma \in \Gamma(G_1)$ and $\gamma' \in \Gamma(\partial F(G_2; p))$ because $u$ is split. Therefore we have that $\beta_\omega(\tilde{f}_p) = \beta_\omega(\tilde{u}_p) = 0$. This completes the proof. □



Figure 6.1.



Figure 6.2.

---

[3] The value of $\overline{\mathrm{Arf}}(L)$ is called the *reduced Arf invariant* of $L$ [15].

**Example 6.2.** Let $G$ be a disjoint union of a circle $\gamma$ and the *handcuff graph* (resp. 2-*bouquet*) $G_2$. Let $\omega$ be a weight on $\Gamma(\gamma)$ over $\mathbb{Z}_2$ defined by $\omega(\gamma) = 1$. We fix an embedding $p : G_2 \rightarrow S^2$ and take a regular neighborhood $F(G_2; p)$ as illustrated in Figure 6.3 (1) (resp. (2)).



Figure 6.3.

Let $f$ be a spatial embedding of $G$ as illustrated in Figure 2.1 (1) (resp. (2)). Let us take an induced embedding $\tilde{f}_p : \gamma \cup F(G_2; p) \rightarrow S^3$ as illustrated in Figure 6.4 (1) (resp. (2)). Note that $\mathrm{lk}(\tilde{f}_p(\gamma), \tilde{f}_p(\gamma')) = 0$ for any $\gamma' \in \Gamma(\partial F(G_2; p))$. Then it can be calculated that $\beta_\omega(\tilde{f}_p) = 1$. Thus by Theorem 6.1 we have that $f$ is non-splittable up to edge-homotopy.



Figure 6.4.

## Acknowledgment

The authors are very grateful to Professor Hitoshi Murakami for his hospitality at the Tokyo Institute of Technology where this work was conducted.

## References

[1] T. D. Cochran, Concordance invariance of coefficients of Conway's link polynomial, *Invent. Math.* **82** (1985), 527–541. MR **87c:**57002

[2] N. Habegger and X. -S. Lin, The classification of links up to link-homotopy, *J. Amer. Math. Soc.* **3** (1990), 389–419. MR **91e:**57015

[3] R. Hartley, The Conway potential function for links, *Comment. Math. Helv.* **58** (1983), 365–378. MR **85h:**57006

[4] F. Hosokawa, On ∇-polynomials of links, *Osaka Math. J.* **10** (1958), 273–282. MR **21**#1606

[5] J. Hoste, The first coefficient of the Conway polynomial, *Proc. Amer. Math. Soc.* **95** (1985), 299–302. MR **86m:**57009

[6] L. H. Kauffman, *Formal knot theory*, Mathematical Notes, **30**, Princeton University Press, Princeton, NJ, 1983. MR **85b:**57006

[7] J. P. Levine, An approach to homotopy classification of links, *Trans. Amer. Math. Soc.* **306** (1988), 361–387. MR **88m:**57008

[8] J. Milnor, Link groups, *Ann. of Math. (2)* **59** (1954), 177–195. MR **17,70e**

[9] T. Motohashi and K. Taniyama, Delta unknotting operation and vertex homotopy of graphs in $R^3$, KNOTS '96 (Tokyo), 185–200, World Sci. Publishing, River Edge, NJ, 1997. MR **99i:**57021

[10] R. Nikkuni, Delta link-homotopy on spatial graphs, *Rev. Mat. Complut.* **15** (2002), 543–570. MR **2004d:**57013

[11] R. Nikkuni, Edge-homotopy classification of spatial complete graphs on four vertices, *J. Knot Theory Ramifications* **13** (2004), 763–777. MR **2005f:**57008

[12] R. Nikkuni, Sharp edge-homotopy on spatial graphs, *Rev. Mat. Complut.* **18** (2005), 181–207. MR2135538

[13] Y. Ohyama and K. Taniyama, Vassiliev invariants of knots in a spatial graph, *Pacific J. Math.* **200** (2001), 191–205. MR **2003a:**57025

[14] N. Sato, Cobordisms of semiboundary links, *Topology Appl.* **18** (1984), 225–234. MR **86d:**57010

[15] T. Shibuya, Self $\sharp$-unknotting operations of links, *Mem. Osaka Inst. Tech. Ser. A* **34** (1989), 9–17. MR **92a:**57014

[16] T. Shibuya and A. Yasuhara, Classification of links up to self pass-move, *J. Math. Soc. Japan* **55** (2003), 939–946. MR **2004f:**57016

[17] R. Sturm Beiss, The Arf and Sato link concordance invariants, *Trans. Amer. Math. Soc.* **322** (1990), 479–491. MR **91m:**57006

[18] K. Taniyama, Link homotopy invariants of graphs in $R^3$, *Rev. Mat. Univ. Complut. Madrid* **7** (1994), 129–144. MR **95f:**57023

[19] K. Taniyama, Cobordism, homotopy and homology of graphs in $R^3$, *Topology* **33** (1994), 509–523. MR **95h:**57002

Department of Mathematics, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

*E-mail address*: `tfleming@math.ucsd.edu`

Institute of Human and Social Sciences, Faculty of Teacher Education, Kanazawa University, Kakuma-machi, Kanazawa, Ishikawa, 920-1192, Japan

*E-mail address*: `nick@ed.kanazawa-u.ac.jp`